



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

First case of human cerebral *Taenia martis* cysticercosis

Brunet, Julie ; Benoild, Aurélien ; Kremer, Stéphane ; Dalvit, Constanza ; Lefebvre, Nicolas ; Hansmann, Yves ; Chenard, Marie-Pierre ; Mathieu, Bruno ; Grimm, Felix ; Deplazes, Peter ; Pfaff, Alexander W ; Abou-Bacar, Ahmed ; Marescaux, Christian ; Candolfi, Ermanno

Abstract: *Taenia martis* is a tapeworm affecting mustelids, with rodents serving as intermediate hosts. The larval stage (cysticercus) has before been found only rarely in humans or primates. We hereby describe a case of cerebral *T. martis* cysticercosis in a French immunocompetent patient, confirmed by DNA analyses of biopsy material.

DOI: <https://doi.org/10.1128/JCM.01033-15>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-111487>

Journal Article

Accepted Version

Originally published at:

Brunet, Julie; Benoild, Aurélien; Kremer, Stéphane; Dalvit, Constanza; Lefebvre, Nicolas; Hansmann, Yves; Chenard, Marie-Pierre; Mathieu, Bruno; Grimm, Felix; Deplazes, Peter; Pfaff, Alexander W; Abou-Bacar, Ahmed; Marescaux, Christian; Candolfi, Ermanno (2015). First case of human cerebral *Taenia martis* cysticercosis. *Journal of Clinical Microbiology*, 53(8):2756-2759.

DOI: <https://doi.org/10.1128/JCM.01033-15>

1 **First case of human cerebral *Taenia martis* cysticercosis**

2

3 Julie BRUNET^{1,2}, Aurélien BENOILID³, Stéphane KREMER⁴, Constanza DALVIT³, Nicolas

4 LEFEBVRE⁵, Yves HANSMANN⁵, Marie-Pierre CHENARD⁶, Bruno MATHIEU², Felix

5 GRIMM⁷, Peter DEPLAZES⁷, Alexander W. PFAFF^{1,2}, Ahmed ABOU-BACAR^{1,2}, Christian

6 MARESCAUX³, and Ermanno CANDOLFI^{1,2}

7 **Authors' affiliations:**

8 Laboratoire de Parasitologie et de Mycologie Médicale, Plateau Technique de Microbiologie,

9 Hôpitaux Universitaires de Strasbourg, F-67091 Strasbourg; ¹ Institut de Parasitologie et Pathologie

10 Tropicale, EA 7292, Fédération de Médecine Translationnelle, Université de Strasbourg, F-67000

11 Strasbourg; ² Service de neurologie, Hôpitaux Universitaires de Strasbourg, F-67098 Strasbourg; ³

12 Département de Neuroradiologie, Hôpitaux Universitaires de Strasbourg, F-67000 Strasbourg; ⁴

13 Service des Maladies Infectieuses et Tropicales, Centre Hospitalier Universitaire, F-67000 Strasbourg;

14 ⁵ Centre hospitalier de Haute-pierre, Laboratoire d'Anatomie Pathologique, F-67000 Strasbourg; ⁶

15 Institute of Parasitology, Vetsuisse Faculty, University of Zurich, Winterthurerstr, 266a CH-8057

16 Zurich⁷

17

18 **Corresponding author:** Julie Brunet

19 Institut de Parasitologie et Pathologie Tropicale, EA 7292, Université de Strasbourg, 3 rue

20 Koeberlé, F-67000 Strasbourg.

21 Tel: + 33 (0)3 68 85 37 00; fax: +33 (0)3 68 85 38 09

22 Email: jbrunet@unistra.fr

23 **Keywords:** *Taenia martis*, cysticercosis, zoonosis

24 **Abstract**

25 *Taenia martis* is a tapeworm affecting mustelids, with rodents serving as intermediate hosts.

26 The larval stage (cysticercus) has before been found only rarely in humans or primates. We
27 hereby describe a case of cerebral *T. martis* cysticercosis in a French immunocompetent
28 patient, confirmed by DNA analyses of biopsy material.

29

30 **Case report**

31

32 In January 2012, a 44-year-old woman with no significant medical history except Verneuil's
33 disease was admitted to the Strasbourg University Hospital for right hemiparesis and aphasia.
34 Cerebral magnetic resonance imaging (MRI) revealed fluid attenuated inversion recovery
35 (FLAIR) and diffusion signal abnormalities in a small area of the left temporal lobe cortex,
36 associated with leptomeningeal gadolinium-based enhancement in the depth of the adjacent
37 sulcus (Fig. 1, 2). On examination, the patient was found to have a temperature of 38.5°C. She
38 reported no travel history outside of Europe and presented no immunosuppression.
39 Meningoencephalitis was suspected and the first examination of the cerebrospinal fluid (CSF)
40 revealed lymphocytic pleocytosis, increased protein levels, and normal glucose concentration.
41 Bacterial antigen detection in CSF was negative, as were CSF bacterial and fungal cultures.
42 Acyclovir, cefotaxime, and amoxicillin were initiated. Five days later, a second lumbar
43 puncture revealed 790 leucocytes/mm³ (70% lymphocytes, 30% eosinophils) and oligoclonal
44 immunoglobulin bands.

45 Due to the eosinophilia, a complementary serological investigation was performed. Serology
46 was weakly positive (1:60) for *Taenia solium*, using *T. solium* antigens and
47 immunofluorescence, and negative for *Echinococcus granulosus* using Western blot with
48 *Echinococcus* spp. antigens (LDBIO Diagnostics, Lyon, France), *E. multilocularis* using
49 enzyme-linked immunosorbent assay (ELISA) with Em2+ antigens, (Bordier Affinity
50 Products SA, Crissier, Switzerland), *Taenia saginata* using *T. saginata* antigens and
51 immunofluorescence, *Schistosoma mansoni* using ELISA with *S. mansoni* antigens (Bordier
52 Affinity Products SA), *Trichinella spiralis* using ELISA with *T. spiralis* antigens (LDBIO
53 Diagnostics), and *Strongyloides stercoralis* using ELISA with *S. ratti* antigens (Bordier

54 Affinity Products SA). Stool examinations were negative for parasites. The patient fully
55 recovered the following week. She was re-admitted in May 2012 after another episode of
56 aphasia. Cerebral MRI was then performed again, displaying a ring-enhancing lesion along
57 with a small enhancing nodule attached to its border, associated with significant perilesional
58 edema, in the same cortical area as that recorded on the first MRI.

59 Despite MRI revealing no evidence of neoangiogenesis, due to the cerebral blood volume
60 being diminished, there were no clinical features that enabled the distinction from an
61 inflammatory process due to either unknown infection or tumor, such as a glioblastoma or
62 metastatic process. Surgical exeresis of the parasitic mass was performed. The pathological
63 examination demonstrated no signs of a tumoral process, yet revealed a thick-walled parasitic
64 cyst with dense fibrosis and intense mononuclear inflammation. The cyst contained dense
65 fluid consisting of thick bright eosinophilic ribbons of membranous tissue and calcareous
66 corpuscles, characteristic of cestodes. Several punch biopsies of the suspected lesion were
67 carried out by means of laser capture microdissection.

68

69 DNA was extracted from the dissected tissues. Polymerase chain reaction (PCR) testing was
70 negative for *Echinococcus granulosus* and *E. multilocularis*, though positive for *Taenia* spp.
71 (1). Fragments of the mitochondrial cytochrome c oxidase subunit 1 (*cox1*), NADH
72 dehydrogenase subunit 1 (*nad1*), and 12S ribosomal ribonucleic acid (rRNA) (*rRNAS*) genes
73 were amplified by means of PCR using primer pairs JB3/JB4.5 (2), JB11/JB12 (3), and
74 Cest3/Cest5 (4), respectively. Both partial sequences of the *cox1* (396bp) and *nad1* (488bp)
75 genes demonstrated 100% homology with published *T. martis* sequences (EU544557 and
76 EU544607). For the rRNA gene, the sequence (263/265bp) demonstrated 99.2% homology
77 with a published *T. martis* sequence (JX415820). Lower sequence identities were also
78 revealed between our specimen and *T. crassiceps* (90 and 82% for *cox1* and *nad1*,

79 respectively) or *T. twitchelli* (94 and 86% for *cox1* and *nad1*, respectively). Cerebral
80 cysticercosis diagnosis was thus established based on the presence of *T. martis*. A
81 phylogenetic tree was created using the sequences obtained in this study as well as those of
82 representative *Taenia* species accessed from GenBank (Fig. 3).

83 Surgical exeresis of the parasitic mass was performed, and the patient was treated with
84 praziquantel (50mg/Kg/day) for 15 days and albendazol (15mg/Kg/day) for 1 month,
85 combined with corticosteroids (1mg/Kg/day). Three years thereafter, no signs of recurrence
86 were observed.

87 The patient lives in the forested part of the French Vosges region, consisting of a woody
88 mountainous area at medium altitude with temperate climate in eastern France. She has three
89 cats and a dog, with a garden where she grows vegetables, with free access to the nearby
90 forest. Fecal samples collected from her cats and dog were investigated and taeniid eggs were
91 detected in one cat and the dog. The eggs were identified as *Taenia taeniaeformis* (4). *T.*
92 *martis* was not identified. Fecal samples collected from the house's surroundings, suspected
93 to originate from martens, were also negative for *T. martis*.

94

95 Several cestode species are currently known to be human parasites. Some are capable of
96 invading the central nervous system (CNS), namely *Taenia solium* and *Echinococcus*
97 *granulosus*. Neurocysticercosis caused by *T. crassiceps* or *T. multiceps* is less common, yet
98 should be considered in the differential diagnosis of CNS infections. *T. martis* cysticercosis
99 was recently reported for the very first time in a human eye (5).

100 *T. martis* is usually found in adult stages in the small intestines of stone martens. Other
101 carnivores, including mustelids of the *Mustela*, *Meles*, *Lutra* or *Gulo* genera, and, more rarely,
102 canids of the *Canis* and *Vulpes* genera, as well as wild cats, may also act as definitive hosts

(6). Rodents and other small mammals serve as intermediate hosts and harbor the parasite in its larval stage (cysticerci). In rodents, cysticerci are usually found in the pleural and peritoneal cavities. The distribution of *T. martis* ranges from Europe (*T. m. martis* subspecies) to North America and Russia (*T. m. americana*) (6). In Europe, *T. martis* has been observed in martens and rodents in Italy, Germany, The Netherlands, Belgium, Spain, Poland, Belarus, and Switzerland (7-9). Recently, fatal *T. martis* cysticercosis was identified in a Tonkean macaque (*Macaca tonkeana*) in Eastern France near Strasbourg (Alsace region) (10) and in a ring-tailed lemur (*Lemur catta*) in the zoological gardens of Rome, Italy (11). These recent cases are evidence of potential lethal outcomes in monkeys and offer particular epidemiological value as indicators of existing environmental contamination with *T. martis* eggs. So far, the only human case that has been described involved the eye of an immunocompetent woman in Germany. The patient exhibited a subretinal tumor-like structure resembling a cestode larva, which was treated with albendazole and dexamethasone, and the cyst was removed. The patient lived in southwestern Germany on the east side of the Rhine River, just across from the Alsace area where our case occurred, and the consumption of contaminated homegrown vegetables was suspected as causative risk factor (5). In our patient, the *T. martis* cysticerci developed in the brain and, once treated, her clinical evolution turned out favorable. Both cases displayed one unique lesion, potentially due to the non-multiplying nature of *T. martis* larvae, thus limiting the risk of generalization and recurrence (6). Transmission to intermediate hosts occurs *via* the oral route by way of food or water contaminated by fecal matter. In our case, the potential risk factors could have been the consumption of wild berries or garden-grown vegetables contaminated by infected marten or fox feces, as the patient lives near the forest and the intrusion of wild animals into the garden is possible. The involvement of her pet animals could not be excluded either, even though the corresponding fecal examinations were negative at the time of sample collection. It is highly

128 probable that a natural *T. martis* life cycle existed in the environment of the house, given that
129 small rodents (voles, mice, rats, and squirrels) and carnivores (martens and foxes) were
130 regularly observed in close vicinity. In France, the prevalence of *T. martis* in wildlife remains
131 largely unknown, with only few reports available (12, 13). However, the recent observation of
132 a peritoneal *T. martis* cysticercosis in a Tonkean macaque in the Alsace region indicates that
133 local contaminations of this particular environment with eggs of this tapeworm species may
134 occur.

135 In summary, we hereby describe the first case of a human cerebral infection with *T. martis*. A
136 treatment comprising praziquantel and albendazole as well as surgical exeresis led to a
137 complete cure. Our study further documents the susceptibility of humans to larval *T. martis*
138 infections. Even though the number of diagnosed cases is low, we cannot rule out the
139 possibility that cases may have been missed in the past due to the absence of molecular
140 analysis. We therefore believe that both *T. martis* and *T. crassiceps* cysticercosis should be
141 added to the differential diagnosis of unusual tumor-like CNS lesions.

142

143 **References**

- 144 1. **Georges, S., O. Villard, D. Filisetti, A. Mathis, L. Marcellin, Y. Hansmann, and**
145 **E. Candolfi.** 2004. Usefulness of PCR analysis for diagnosis of alveolar
146 echinococcosis with unusual localizations: two case studies. *J. Clin. Microbiol.*
147 **42**:5954-5956.
- 148 2. **Bowles, J., D. Blair, and D.P. McManus.** 1992. Genetic variants within the genus
149 *Echinococcus* identified by mitochondrial DNA sequencing. *Mol. Biochem. Parasitol.*
150 **54**:165-173.
- 151 3. **Bowles, J., and D.P. McManus.** 1993. NADH dehydrogenase 1 gene sequences
152 compared for species and strains of the genus *Echinococcus*. *Int. J. Parasitol.* **23**:969-
153 972.
- 154 4. **Trachsel, D., P. Deplazes, and A. Mathis.** 2007. Identification of taeniid eggs in the
155 faeces from carnivores based on multiplex PCR using targets in mitochondrial DNA.
156 *Parasitology.* **134**:911-920.
- 157 5. **Eberwei, P., A. Haeupler, F. Kuepper, D. Wagner, W.V. Kern, B. Muntau, P.**
158 **Racz, H. Agostini, and S. Poppert.** 2013. Human infection with marten tapeworm.
159 *Emerg. Infect. Dis.* **19**:1152-1154.
- 160 6. **Loos-Frank, B.** 2000. An up-date of Verster's (1969) 'Taxonomic revision of the
161 genus *Taenia Linnaeus*' (Cestoda) in table format. *Syst. Parasitol.* **45**:155-183.
- 162 7. **Ribas, A., I. Torre, C. Feliu, A. Arrizabalaga, and J.C. Casanova.** 2009. Helminth
163 communities of the bank vole *Myodes glareolus* (Rodentia, Arvicolinae) in two
164 populations: Montsenyu Natural Park (north-eastern-Spain) and Pi Natural Reserve
165 (French Pyrenees). *Rev. Ibero-latinoam. Parasitol. Int.* **1**:73-81.

- 166 8. **Mathy, A., R. Hanosset, S. Adant, and B. Losson.** 2009. The carriage of larval
167 *Echinococcus multilocularis* and other cestodes by the musk rat (*Ondatra zibethicus*)
168 along the Ourthe River and its tributaries (Belgium). *J. Wildl. Dis.* **45**:279-287.
- 169 9. **Reperant, L.A., D. Hegglin, I. Tanner, C. Fischer, and P. Deplazes.** 2009. Rodents
170 as shared indicators for zoonotic parasites of carnivores in urban environments.
171 *Parasitology.* **136**:329-337.
- 172 10. **Brunet, J., B. Pesson, R. Chermette, P. Regnard, F. Grimm, P. Deplazes, X.**
173 **Ferreira, M. Sabou, A.W. Pfaff, A. Abou-Bacar, and E. Candolfi.** 2014. First case
174 of peritoneal cysticercosis in a non-human primate host (*Macaca tonkeana*) due to
175 *Taenia martis*. *Parasit. Vectors.* **7**:422.
- 176 11. **De Liberato, C., F. Berrilli, R. Meoli, K.G. Friedrich, P. Di Cerbo, C. Cocumelli,**
177 **and C. Eleni.** 2014. Fatal infection with *Taenia martis* metacestodes in a ring-tailed
178 lemur (*Lemur catta*) living in an Italian zoological garden. *Parasitol. Int.* **63**:695-7.
- 179 12. **Joyeux, C., and J.G. Bear.** 1934. Sur quelques Cestodes de France. *Archives du*
180 *Muséum National d'Histoire Naturelle.* **sér.6**:157-171.
- 181 13. **Ribas, A., C. Milazzo, P. Foronda, and J.C. Casanova.** 2004. New data on
182 helminths of stone marten, *Martes foina* (Carnivora, Mustelidae), in Italy.
183 *Helminthologia.* **41**:59-61.
- 184 14. **Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar.** 2011.
185 MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood,
186 Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.* **28**: 2731-
187 9.
- 188

189 **FIGURE LEGENDS**

190

191 **FIG. 1.** Axial fluid attenuated inversion recovery (FLAIR) (a, e); axial diffusion (b); apparent
192 diffusion coefficient map (c); axial T1 Spin Echo (SE) after gadolinium injection (d). Mild
193 cortical FLAIR and diffusion hypersignal, with mild restriction of the diffusion coefficient in
194 the left temporal lobe (arrow). Leptomeningeal enhancement after gadolinium administration
195 in the depth of the adjacent sulcus (arrow head). White matter subcortical nonspecific
196 hyperintensities (star)

197

198 **FIG. 2.** Sagittal T1 (a); axial T2 Spin Echo (SE) (b); T2 star-weighted angiography (SWAN)
199 (c); axial (d) and sagittal (e) T1 SE after gadolinium injection; axial diffusion (f); apparent
200 diffusion coefficient map (g); cerebral blood volume map (h). Left temporal cortical annular
201 enhancing lesion (arrow). The border of the lesion is thick and partially spontaneously
202 hyperintense on T1 and hypointense on T2, with T2 SWAN suggesting the presence of
203 hemosiderin or calcium deposits (arrow head). The contrast enhancement is predominant in
204 the inner part of the border (arrow). The center of the lesion appears hyperintense on
205 diffusion-weighted MR imaging, with restriction of the diffusion coefficient suggesting thick
206 content (star). In the center of the lesion, there is a small enhancing nodule attached to the
207 border, hypointense on T2 SWAN (circle). The cerebral blood volume is diminished (h).
208 Significant perilesional edema is revealed

209

210 **FIG. 3.** Neighbor-joining tree based on *CoxI* sequences, showing the phylogenetic
211 relationships between the *Taenia martis* isolated from humans and the other tapeworm

212 species. Shaded triangle refers to the sequence generated by this study and the Genbank
213 accession numbers of each strain are given in parentheses. (a) to (d) refer to *Taenia martis*
214 isolated from humans, *Myodes glareolus*, *Apodemus sylvaticus*, and *Myodes rufocanus*,
215 respectively. Bootstrap values >70% are included to indicate branch support. Neighbor-
216 joining analysis (NJ) was processed using MEGA V.5.2. software and the branch support was
217 evaluated by means of bootstrap test with 1000 replicates (14)

218

219

220





